

INVITRO FERTILIZATION OF BUFFALO OOCYTES CULTURED IN THREE DIFFERENT MEDIA

Bhoopendra Singh Chauhan¹, Ramjanam Dubey²

¹ Research Scholar, Himalayan University, Arunachal Pradesh

² Assistant Professor, D.D. Degree College, Azamgarh
India

Abstract—In order to evaluate the effect of media on in vitro maturation (IVM; 24 hours) and subsequent fertilization (IVF; 24 hours), culturable grade buffalo oocytes (n=1058) were matured (Expt. 1, n=517) or matured and fertilized (Expt. 2, n=541) in vitro in three different media (Ham's F-10, Waymouth MB and TCM-199) with same supplements (5µg/ml FSH, 5µg/ml LH, 1µg/ml estradiol, 25mMHepes, 0.25mMpyruvate and antibiotics) to record two end points; IVM and IVF. The overall mean culturable grade oocyte recovery was 3.12±0.20. At the end of Expt. 1 significantly higher (P<0.01) proportion of oocytes were matured in vitro in Waymouth compared to Ham's F-10. In TCM-199, IVM rates were non-significantly higher compared to Ham's F-10 and non-significantly lower compared to Waymouth MB medium. At the end of Expt. 2 the proportion of oocytes that fertilized were 10%, 17% and 16.4% in Ham's F-10, Waymouth and TCM-199 respectively which were non significantly different. It was concluded that Way mouth MB medium is the most appropriate medium for in vitro maturation of buffalo oocytes followed by TCM-199 and Ham's F 10 however the IVM media had a negligible effect on the subsequent fertilization of buffalo oocytes.

Index Terms— Buffalo, In vitro maturation, In vitro fertilization, Media, Oocytes.

I. INTRODUCTION

Buffalo oocyte IVM rates were low in initial studies, but improved with the addition of buffalo serum or hormones (Hammam et al., 2010) in the culture medium. Addition of cumulus cells alone did not improve the oocyte maturation rate (Das et al., 1997). Expensive components of in vitro maturation (IVM) medium, such as fetal calf serum and hormones, were successfully replaced by steer serum and follicular fluid (Nandi et al., 2002). The effect of supplements in the medium such as cysteamine, growth factors and PMSG was considered useful to promote in vitro maturation and subsequent fertilization of buffalo oocytes. A large number of variables that influence in vitro maturation of buffalo oocytes such as follicular size from which the oocytes are recovered, oocyte recovery procedures, presence or absence of cumulus cells, supplements in the media have been explained and recently (Mahmoud and El-Naby, 2013). In spite of several attempts to improve the in vitro

maturation and subsequent fertilization of buffalo oocytes the results obtained had been modest and inconsistent (Hegab et al., 2009). The composition of media appears to be an important regulator of sequential in vitro maturation and subsequent development of follicular oocytes (Hammam et al., 2010). The commonly used media for IVM of buffalo oocytes is TCM 199 (Deneke et al., 2013) however, since this media is complex other culturable media such as Ham's F 10 (Hammam et al., 2010) DMEM (Hegab et al., 2009) Ferti Cult medium (Hegabet al., 2009), Ham's F 12 have been experimented with variable results. The use of defined synthetic medium such as Way mouth MB medium have shown some advantage over TCM 199 when supplemented with growth factors (Purohit et al., 2005). The in vitro maturation and subsequent development of bubaline oocytes in different media with the same supplements is sparsely studied. In this study we evaluated TCM- 199, Waymouth MB medium and Hams F-10 for in vitro maturation and subsequent fertilization of buffalo follicular oocytes.

II. MATERIALS AND METHODS

Buffalo ovaries (n=368) were collected from a local abattoir in warm normal saline and brought to the laboratory. Surface follicles from the ovaries were aspirated to collect the oocytes. The culturable grade oocytes (n=1058) were randomly allocated to three different media (Ham's F-10, Waymouth MB and TCM-199) to record two end points in vitro maturation (Experiment 1) and in vitro fertilization (Experiment 2) in two separate experiments. In experiment 1, all oocytes that were matured for 24 hours were fixed and evaluated for nuclear maturation (proportion reaching Metaphase- II). In experiment 2 all oocytes were first matured in vitro and after 24 hours were fertilized with prepared buffalo spermatozoa. After 24 hours of sperm oocyte co-incubation the presumptive zygotes were fixed and evaluated for in vitro fertilization. All media and chemicals were from Sigma chemical company USA.

The oocytes were divided into three groups of approximately equal number of oocytes and cultured in either TCM-199, Waymouth MB media or Hams F-10 with the same

supplements (5µg/ml FSH, 5µg/ml LH, 1µg/ml estradiol, 25mMHepes, 0.25mM Pyruvate and antibiotics) in 50-100µl maturation media covered by sterile paraffin oil for 24 hours at 38±1°C and 5%CO₂ in humidified air in a CO₂ incubator for in vitro maturation. After 24 hours of maturation, all oocytes from different groups were collected and fixed separately for staining. The surrounding cumulus cells were removed by vortexing for 1 minute in TCM 199 with hyaluronidase (0.3%). The oocytes were placed in the center of an area delineated by two paraffin wax bars on a clean grease free glass slide. The denuded oocytes were compressed gently with a cover slip to hold and were fixed for 24 hours in acetic acid and methanol [1:3(v: v)] and stained with 1% Giemsa stain for evaluation of nuclear status. Oocytes were considered mature if they were at metaphase-II.

Oocytes were matured in vitro in the three different media utilizing the same procedures as above, and after 24 hours of maturation they were fertilized with prepared buffalo spermatozoa.

Frozen thawed buffalo semen was prepared for IVF using a discontinuous Percoll density gradient to separate highly motile live spermatozoa as per Grant et al. (1994). Briefly 4 ml of 90% isotonic Percoll was layered in a 15 ml centrifuge tube beneath 4 ml of 40% isotonic Percoll. The sperms were washed initially in TALP-BSA by centrifugation at 250 g for

4 minutes. The sperm pellet was resuspended in 1 ml of the medium. The washed sperm pellet was layered on the top of Percoll gradient and centrifuged at 300 g for 35 min. The resultant pellet was removed from the bottom and washed twice in TALP-BSA by centrifugation.

The sperm pellet was re-suspended in TALP to give a final concentration of 1-2 million sperms. This was incubated for 2-3 hours in a CO₂ incubator. The matured oocytes were transferred to another dish containing Fert-TALP medium (TALP supplemented with 30 µg/ml penicillamine, 15 µmol/ml hypotaurine, 10 µg/ml Heparin and 1 µmol/ml adrenaline) under paraffin oil. They were inseminated with prepared sperms in a volume, so as to give a final concentration of 1-2 million sperms.

Following co-incubation for 20-24 hours with sperms, all the oocytes from each group were washed with fresh medium and vortexed for 1 minute to separate the cumulus mass. They were processed for fixing and staining in the same way as oocytes were fixed after IVM. Oocytes were considered fertilized if they revealed 2 pronuclei or a swollen sperm head along with M-II plate as described previously (Purohit et al., 2005). The arcsine transformed data of the proportion of oocytes matured or fertilized was compared by one way ANOVA.

III. RESULTS AND DISCUSSION

The overall mean number of oocytes recovered per ovary was 3.66 ± 0.24 whereas the overall mean number of culturable oocyte recovery per ovary was 3.12 ± 0.20. Similar recovery rates were observed in a few studies on buffalo

(Mistry and Dhama, 2009). However a large number of previous studies had recorded a lower culturable grade oocyte recovery rates varying from 0.4-2.17. The reasons for differences in the oocyte recovery rates are diverse and include reproductive status of the animal from which they are retrieved, presence or absence of CL, season of recovery and recovery procedure adopted (Mehmood et al., 2011).

In vitro maturation of buffalo oocytes in Expt 1 and Expt 2 revealed that significantly higher (P<0.01) proportion of oocytes matured in vitro (reached M-II stage) in Way mouth MB media compared to Ham's F 10 media. The number of oocytes that matured in vitro was non-significantly higher (P>0.01) in Way mouth MB media compared to TCM 199 suggesting better performance of oocytes in Way mouth MB media. The overall maturation rates (all three media) obtained during the present study were 70.01 percent. Similar maturation rates were recorded in many previous studies on buffalo oocytes matured in vitro (Leal et al., 2010).

Previous studies on buffalo oocyte maturation in vitro have shown TCM-199 to be better over Hams F-10 (Hammam et al., 2010). The beneficial effect of TCM-199 on IVM may be related to some factors in its composition such as essential amino acids and glutamine that stimulate DNA and RNA synthesis and enhance cell division (Mahmoud and El-Naby, 2013).

Waymouth medium was found to support in vitro maturation of buffalo oocytes even better to TCM-199. Xu et al. (1992) have previously shown that Waymouth medium yielded better cleavage rates compared to TCM-199 during bovine in vitro embryo development. A previous study on buffalo oocytes (Purohit et al., 2005) had recorded comparable in vitro maturation rates both with TCM-199 and Waymouth medium.

In Expt 2, in vitro maturation in Way mouth MB media resulted in subsequent higher fertilization rates (17%) compared to Ham's F 10 (10%) and TCM 199 (16.4%) however the differences were non-significant.

The fertilization rates recorded during the present study are similar to a few of previous study that recorded fertilization rates varying from 11-24% for buffalo oocytes (Hammam et al., 2010). However many other studies recorded higher in vitro fertilization rates (43-82%) for buffalo oocytes (Mehmood et al., 2011). This could be because of the difference in the supplements used, the initial quality of oocytes utilized and other variables.

Although differences in the in vitro fertilization rates for buffalo oocytes across various media used for their in vitro maturation have been recorded (Kumar et al., 2008) however it is a general consensus based on analysis of many reports that the supplements are much more important and concur with the findings of the present study.

Comparison of in vitro maturation and fertilization of buffalo oocytes across the three media revealed that in vitro maturation was better in Waymouth and TCM media compared to Hams F-10 in both experiment 1 and experiment 2 and although the fertilization rates were also higher yet the

differences across media are more operative during in vitro maturation of oocytes. It was concluded that Way mouth MB medium is the most appropriate medium for in vitro maturation of buffalo oocytes followed by TCM-199 and Ham's F 10 however, the IVM media had a negligible effect on the subsequent fertilization of buffalo oocytes.

REFERENCES

- [1] Das, S.K., Chauhan, M.S., Palta, P., Tomer, O.S. (1997). Influence of cumulus cells on in vitro maturation of denuded buffalo oocytes. *Vet. Rec.*, 141:522-523.
- [2] Deneke, Y., Yadav, P.S., Deb, R. and Nanda, T. (2013). Comparative studies of the effect of BSA Vs FCS as a supplement in TCM-199 on the in vitro maturation rate of buffalo oocytes collected from slaughter house ovaries. *Buffalo Bull.*, 32:21-25.
- [3] Drost, M. (2007). Advanced reproductive technology in the water buffalo. *Theriogenology*, 68:450- 453.
- [4] Grant, S.A., Long, S.E. and Parkinson, T.J. (1994). Fertilizability and structural properties of boar spermatozoa prepared by Percoll gradient centrifugation. *J. Reprod. Fertil.*, 100:477-483.
- [5] Hammam, A.M., Whisnant, C.S., Elias, A., Zaabel, S.M., Hegab, A.O. and Abu-El-Naga, E.M. (2010). Effect of media, sera and hormones on in vitro maturation and fertilization of water buffalos (*Bubalus bubalis*). *J. Anim. Vet. Adv.*, 9:27-31.
- [6] Hegab, A.O., Montasser, A.E., Hammam, A.M., El-Naga, E.M.A.A. and Zaabel, S.M. (2009). Improving in vitro maturation and cleavage rates of buffalo oocytes. *Anim. Reprod.*, 6:416-421.
- [7] Kumar, D., Anand, T. and Chauhan, M.S. (2008). In vitro development of buffalo embryos using simple media. *Indian Vet. J.*, 85:819-821.
- [8] Leal, L.S., Moya-Araujo, C.F., Fernandes, C.B., Martins, L.R., Landim-Alvarenga, F.C. and Oba,
- [9] E. (2010). Evaluation of recovery, quality and in vitro nuclear maturation of oocytes obtained from buffalo and bovine ovaries. *Revista Vet.*, 21:892-894.
- [10] Madan, M.L., Singla, S.K., Jaiikhani, S. and Ambrose, J.D. (1991). In vitro fertilization in buffalo and birth of first ever IVF buffalo calf. In *Proc. Third World Buffalo Congr. Varma, Bulgaria*, 7:11-17.
- [11] Mahmoud, K.G.M. and El-Naby, A.H.H. (2013). Factors affecting buffalo oocyte maturation. *Global Vet.*, 11:497-510.
- [12] Mehmood, A., Anwar, M., Andrabi, S.M.H., Afzal, M. and Naqvi, S.M.K. (2011). In vitro maturation and fertilization of buffalo oocytes: the effect of recovery and maturation methods. *Turkish J. Vet. Anim. Sci.*, 35:381-386.
- [13] Mistry, C.N. and Dhama, A.J. (2009). Studies on follicular size and oocytes recovery rate from buffalo ovaries by slicing method. *Indian J. Field Vet.*, 5:23-26.
- [14] Nandi, S., Ravindranatha, B.M., Gupta, P.S.P. and Sarma, P.V. (2002). Timing of sequential changes in cumulus cells and first polar body extrusion during in vitro maturation of buffalo oocytes. *Theriogenology*, 57:1151-1159.
- [15] Palanisamy, A., Rangasamy, S., SatheshKumar, S. and Kumanan, K. (2009). Effect of cysteamine supplementation in semi defined media on in vitro production of buffalo embryos. *Indian J. Anim. Sci.*, 30:30-36.
- [16] Purohit, G.N., Brady, M.S. and Sharma, S.S. (2005). Influence of epidermal growth factor and insulin like growth factor 1 on nuclear maturation and fertilization of buffalo cumulus oocyte complexes in serum free media and their subsequent development in vitro. *Anim. Reprod. Sci.*, 87:229-239.
- [17] Xu, K.P., Yadav, B.R., Rorie, R.W., Plante, L., Betteridge, K.J. and King, W.A. (1992). Development and viability of bovine embryos derived from oocytes matured and fertilized in vitro and co-cultured with bovine oviductal epithelial cells. *J. Reprod. Fertil.*, 94:33-43.